



ACTIVITY OF PRIMARY MEDICAL CARE PATIENTS IN VACCINATION AGAINST INFLUENZA, TICK- BORNE ENCEPHALITIS, HEPATITIS AND PNEUMOCOCCAL DISEASE

Beatričė Aleksaitė, Jomantė Mačiukaitienė, Erika Lingytė

Lithuanian university of health sciences, faculty of medicine

ABSTRACT

Introduction: Infectious diseases do harm not only to human health, but also cause certain consequences for national economics as well. The programme on immunisation is one of the most acknowledged and successful public health programmes in the whole world

Aim: Assessing the activity of primary medical care patients in vaccination against influenza, tick-borne encephalitis, hepatitis and pneumococcal disease.

Methodology: The research was carried out from April 2015 up to March 2016. The data was collected at medical centre *Saulės šeimos medicinos centras* (SŠMC) of Kaunas city. During patients' visits, family doctors filled in anonymous questionnaires about the patients visiting SŠMC for consultation by indicating their gender, age and vaccination. There were 10 family doctors in total from SŠMC taking part in the research.

Results: The research involved 1686 patients: 1022 (60.6%) females and 664 (39.4%) males. The average age of the investigated population was 39.16 (\pm SN) \pm 27.62 of 0 to 94 years of age. The group of randomly-selected patients of 0-18 years of age included 497 persons (29.5%), 19-35 years of age – 280 (16.6%), 36-50 years of age – 260 (15.4%), 51-70 years of age – 358 (21.2%), \geq 71 years of age – 291 (17.3%). The total number of the patients vaccinated with any vaccine amounted to 864 (51.25%), the number vaccinated against influenza was 494 (29.30%), against pneumococcal disease – to 330 (19.60%), against TBE – to 154 (9.10%), against hepatitis – to 130 (7.70%). An influenza vaccine was administered in 331 (32.89%) females and 163 (24.55%) males, $p=0.001$. A pneumococcal vaccine was administered in 199 (19.47%) females and 131 (19.73%) males, $p=0.90$. A TBE vaccine was administered in 97 (9.49%) females and 57 (8.58%) males, $p=0.53$.

Conclusions: 1. The most part of the patients has vaccinated against influenza and pneumococcal disease, the least part – against TBE and hepatitis. 2. The influenza vaccine has been administered in females more often than in males. 3. Vaccines against influenza and pneumococcal disease are mostly used by patients older than 71 years of age. Vaccines against hepatitis and TBE are mostly administered in the patients of 0 to 18 years of age. 4. Most patients have been vaccinated against influenza only. And very little part of the patients has been vaccinated with two or more different vaccines.

PIRMINĖS SVEIKATOS PRIEŽIŪROS PACIENTŲ AKTYVUMAS SKIEPIJANTIS GRIPO, ERKINIO ENCEFALITO, HEPATITO IR PNEUMOKOKO VAKCINOMIS

Beatričė Aleksaitė, Jomantė Mačiukaitienė, Erika Lingytė

LSMU Medicinos fakultetas

Ižanga

Infekcinės ligos daro žalą ne tik žmonių sveikatai, bet turi pasekmių ir šalies ekonomikai. Imunizacijos programa yra viena labiausiai pripažintų ir sėkmingų visuomenės sveikatos programų visame pasaulyje. Ši programa pasiekė reikšmingų rezultatų daugelyje šalių, todėl skiepais gerinama užkrečiamųjų ligų kontrolė. Ypač svarbu nepamiršti imunoprofilaktikos suaugusiems, esantiems rizikos grupėse, kuriems imunizacija pagerintų ligų išėjimą.

Tikslas

Įvertinti pirminės sveikatos priežiūros pacientų aktyvumą skiepijantis gripo, erkinio encefalito, hepatito ir pneumokoko vakcinomis.

Uždaviniai

1. Nustatyti pacientų aktyvumą skiepijantis gripo, erkinio encefalito (EE), hepatito ir pneumokoko vakcinomis.
2. Palyginti besikreipiančių vyrų ir moterų skiepijimosi dažnį.
3. Išsiaiškinti, kokiomis vakcinomis dažniausiai skiepijasi skirtingų amžiaus grupių pacientai.
4. Nustatyti skiepijimosi gripo, EE, hepatito ir pneumokoko vakcinomis dėsningumus.

Metodika


Tyrimas vykdytas nuo 2015 m. balandžio mėn. iki 2016 m. kovo mėn. Duomenys rinkti Kauno miesto Saulės šeimos medicinos centre (SŠMC). Šeimos gydytojai pacientų vizitų metu pildė anonimines anketas apie į SŠMC besikreipiančius pacientus, pažymėdami jų lytį, amžių ir skiepus. Viso tyrime dalyvavo 10 SŠMC šeimos gydytojų, kurie surinko duomenis apie atsitiktinai atrinktus 1686 besikreipiančius pacientus. Analizuojant duomenis pacientai buvo suskirstyti į amžiaus grupes: 0-18, 19-35, 36-50, 51-70 ir ≥ 71 metų grupės. Statistinė duomenų analizė atlikta SPSS 20.0 statistiniu paketu. Duomenys vertinti kaip statistiškai reikšmingi kai $p < 0,05$.

Rezultatai

Į tyrimą buvo įtraukti 1686 pacientai: 1022 (60,6 proc.) moterys ir 664 (39,4 proc.) vyrai. Tiriamųjų amžiaus vidurkis $39,16 (\pm SN) \pm 27,62$ nuo 0 iki 94 metų. Iš visų atsitiktinai atrinktų pacientų 0-18 metų amžiaus grupėje buvo 497 tiriamieji (29,5 proc.), 19-35 metų - 280 (16,6 proc.), 36-50 metų - 260 (15,4 proc.), 51-70 metų - 358 (21,2 proc.), ≥ 71 metų - 291 (17,3 proc.).

Viso buvo pasiskiepiję kuria nors vakcina 864 (51,25 proc.) pacientai, gripo vakcina - 494 (29,30 proc.), pneumokoko - 330 (19,60 proc.), EE - 154 (9,10 proc.), hepatito - 130 (7,70 proc.) (1 pav).

PACIENTŲ AKTYVUMAS SKIEPIJANTIS GRIPO, ERKINIO ENCEFALITO, HEPATITO IR PNEUMOKOKO VAKCINOMIS




1 pav. Pacientų aktyvumas skiepijantis gripo, EE, hepatito ir pneumokoko vakcinomis

Gripo vakcina skiepijosi 331 (32,89 proc.) moteris ir 163 (24,55 proc.) vyrai, $p=0,001$. Pneumokoko vakcina skiepijosi 199 (19,47 proc.) moterys ir 131 (19,73 proc.) vyrai, $p=0,90$. EE vakcina skiepijosi 97 (9,49 proc.)

moterys ir 57 (8,58proc.) vyrai, $p=0,53$. Hepatito vakcina skiepijosi 72 (7,05proc.) moterys ir 58 (8,73proc.) vyrai, $p=0,20$ (2 pav.).

VYRŲ IR MOTERŲ SKIEPIJIMOSI DAŽNIS




2 pav. Vyrų ir moterų skiepijimosi dažnių palyginimas

Gripo vakcina <18 amžiaus grupėje skiepijosi 50 (10,06proc.) pacientų, 19-35 grupėje – 92 (32,86proc.), 36-50 grupėje – 71 (27,30proc.), 51-70 grupėje – 144 (40,22proc.), ≥ 71 grupėje – 137 (47,08proc.) pacientai, p=0,0001. Pneumokoko vakcina <18 amžiaus grupėje skiepijosi 111 (22,33proc.) pacientų, 19-35 grupėje – 42 (15,00proc.), 36-50 grupėje – 33 (12,69proc.), 51-70 grupėje – 73 (20,39proc.), ≥ 71 grupėje – 71 (24,40proc.) pacientas, p=0,001. EE vakcina <18 amžiaus grupėje

skiepijosi 46 (9,66proc.) pacientai, 19-35 grupėje – 27 (6,64proc.), 36-50 grupėje – 18 (6,92proc.), 51-70 grupėje – 31 (8,66proc.), ≥ 71 grupėje – 32 (11,00proc.) pacientai, p=0,57. Hepatito vakcina <18 amžiaus grupėje skiepijosi 120 (24,14proc.) pacientų, 19-35 grupėje – 7 (2,5proc.), 36-50 grupėje – 2 (0,77proc.), 51-70 grupėje – 1 (0,28proc.) ≥ 71 grupėje pasiskiepijusių nebuvo, p=0,0001 (3 pav.).

3 pav. Skiepijimosi ypatumai skirtingo amžiaus grupės



Tik gripo vakcina skiepijosi 339 (20,10proc.) pacientai, pneumokoko – 161 (9,50proc.), EE– 74 (4,40proc.), hepatito - 71 (4,20proc.), gripo ir pneumokoko vakcinomis - 98 (5,80proc.) pacientai, gripo ir EE - 31 (1,80proc.) pacientas, gripo ir hepatito - 32 (1,90proc.) pacientai, EE ir hepatito - 7 (0,40 proc.) pacientai, gripo, pneumokoko ir EE vakcinomis - 13 (0,80proc.) pacientų, gripo, pneumokoko ir hepatito - 2 (0,10proc.) pacientai, pneumokoko, EE ir hepatito - 8 (0,50proc.) pacientai, gripo, EE ir hepatito - 1 (0,10proc.) pacientas.

Išvados

1. Daugiausia pacientų skiepijosi gripo ir pneumokoko vakcinomis, mažiausia dalis pacientų skiepijosi EE ir hepatito vakcinomis.
2. Gripo vakcina dažniau skiepijasi moterys nei vyrai.
3. Gripo ir pneumokoko vakcina dažniausiai skiepijasi vyresni nei 71 metų pacientai. Hepatito ir EE vakcina dažniausiai skiepijasi pacientai nuo 0 iki 18 metų.
4. Daugiausia pacientų buvo pasiskiepiję tik gripo vakcina. Ir labai nedidelė pacientų dalis buvo pasiskiepiję dviem ar daugiau skirtingų vakcinų.

Literatūros sąrašas

1. Prof. A. Ambrozaitis, doc. L. Jančorienė, prof. A. Mickienė, doc. D. Velyvytė. Suaugusiųjų rizikos grupių skiepijimo Lietuvoje rekomendacijos 2015.
2. Hill AB., Kilgore C., McGlynn M., Jones CH. Improving global vaccine accessibility. 2016 Mar 16;42:67-73.
3. Lahariya C. Health system approach" for improving immunization program performance. 2015 Oct-Dec;4(4):487-94.
4. Nowalk MP, Zimmerman RK, Lin CJ, Reis EC, Huang HH, Moehling KK, et al. Maintenance of Increased Childhood Influenza Vaccination Rates 1 Year After an Intervention in Primary Care Practices. *Academic pediatrics*. 2016;16(1):57-63.
5. Ngoh HL, Ng MC. Vaccination in the primary care setting: when is it safe to proceed? *Singapore medical journal*. 2016;57(1):3-7.
6. Malo TL, Perkins RB, Lee JH, Vadaparampil ST. Primary Care Physicians' Adherence to Expert Recommendations for Cervical Cancer Screening and Prevention in the Context of Human Papillomavirus

- Vaccination. *Sexually transmitted diseases*. 2016;43(7):438-44.
5. Kaasenbrood F, Hollander M, Rutten FH, Gerhards LJ, Hoes AW, Tieleman RG. Yield of screening for atrial fibrillation in primary care with a hand-held, single-lead electrocardiogram device during influenza vaccination. *Europace : European pacing, arrhythmias, and cardiac electrophysiology : journal of the working groups on cardiac pacing, arrhythmias, and cardiac cellular electrophysiology of the European Society of Cardiology*. 2016;18(10):1514-20.
 6. Hoffmann K, Paget J, Wojczewski S, Katic M, Maier M, Soldo D. Influenza vaccination prevalence and demographic factors of patients and GPs in primary care in Austria and Croatia: a cross-sectional comparative study in the framework of the APRES project. *European journal of public health*. 2016;26(3):395-401.
 7. Glisser MB, Barragan Tdel C, Weisstaub G. [Breastfeeding indicators produced at time of vaccination in four Primary Care Centres in southern Santiago, Chile]. *Revista chilena de pediatria*. 2016;87(1):11-7.
 8. Evans AM, Wood FC, Carter B. National community pharmacy NHS influenza vaccination service in Wales: a primary care mixed methods study. *The British journal of general practice : the journal of the Royal College of General Practitioners*. 2016;66(645):e248-57.
 9. Askelson NM, Edmonds SW, Momany ET, Tegegne MA. Understanding Clinic Practices for Human Papilloma Virus Vaccination Series Completion in Clinics That Provide Primary Care: Survey of Clinic Managers in Iowa. *Sexually transmitted diseases*. 2016;43(7):445-9.
 10. Anton F, Richart MJ, Serrano S, Martinez AM, Pruteanu DF. [Strategies to improve influenza vaccination coverage in Primary Health Care]. *Semergen*. 2016;42(3):147-51.
 11. Acevedo G, Lopez L, Willington A, Burrone S, Farias A, Sanchez J. Association of vaccination coverage with sociodemographic factors in workers of primary health care centers of Cordoba, Argentina. *Revista de la Facultad de Ciencias Medicas*. 2016;73(3):163-9.
 12. Rivera Casares F, Pineda Coronel R, Garcia de Leon Chocano S, Lopez S, Montagud S, Lorente P. [Vaccination status of immunocompromised patients attended in primary care]. *Gaceta sanitaria*. 2015;29(3):235-6.
 13. Ngamruengphong S, Horsley-Silva JL, Hines SL, Pungpapong S, Patel TC, Keaveny AP. Educational

- Intervention in Primary Care Residents' Knowledge and Performance of Hepatitis B Vaccination in Patients with Diabetes Mellitus. *Southern medical journal*. 2015;108(9):510-5.
14. McCrossan P, McCafferty C, Murphy C, Murphy J. Retrospective review of administration of childhood primary vaccination schedule in an Irish tertiary neonatal intensive care unit. *Public health*. 2015;129(7):896-8.
 15. Kunin M, Engelhard D, Thomas S, Ashworth M, Piterman L. Challenges of the Pandemic Response in Primary Care during Pre-Vaccination Period: A Qualitative Study. *Israel journal of health policy research*. 2015;4:32.
 16. Katz A, Enns JE, Chateau D, Lix L, Jutte D, Edwards J, et al. Does a pay-for-performance program for primary care physicians alleviate health inequity in childhood vaccination rates? *International journal for equity in health*. 2015;14:114.
 17. Fu LY, Zook K, Gingold J, Gillespie CW, Briccetti C, Cora-Bramble D, et al. Frequent vaccination missed opportunities at primary care encounters contribute to underimmunization. *The Journal of pediatrics*. 2015;166(2):412-7.
 18. Djibo DA, Peddecord KM, Wang W, Ralston K, Sawyer MH. Factors Associated With Missed Opportunities for Influenza Vaccination: Review of Medical Records in a Diverse Sample of Primary Care Clinics, San Diego County, 2010-2011. *Journal of primary care & community health*. 2015;6(3):147-53.
 19. Arnould B, Arnould P, Benmedjahed K, Coindard G, Denis F, Duhot D, et al. Scoring and Measurement Properties of a Tool to Assess Primary Care Physicians' Engagement in and Perceived Barriers to Vaccination: the 'Determinant of Intentions of Vaccination' (Diva(c)) Questionnaire. *Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research*. 2015;18(7):A596.
 20. Zimmerman RK, Nowalk MP, Lin CJ, Hannibal K, Moehling KK, Huang HH, et al. Cluster randomized trial of a toolkit and early vaccine delivery to improve childhood influenza vaccination rates in primary care. *Vaccine*. 2014;32(29):3656-63.
 21. Yousafzai MT, Qasim R, Khalil R, Kakakhel MF, Rehman SU. Hepatitis B vaccination among primary health care workers in Northwest Pakistan. *International journal of health sciences*. 2014;8(1):67-76.
 22. William S, Reid J. IPCRG:International Primary Care Respiratory Group IPCRG World Conference, International Research Collaboration, Asthma Resources and Motivation for Flu Vaccination. *The European journal of general practice*. 2014;20(1):85-6.
 23. Weltermann BM, Markic M, Thielmann A, Gesenhues S, Hermann M. Vaccination management and vaccination errors: a representative online-survey among primary care physicians. *PloS one*. 2014;9(8):e105119.
 24. Wang K, Fry NK, Campbell H, Amirthalingam G, Harrison TG, Mant D, et al. Whooping cough in school age children presenting with persistent cough in UK primary care after introduction of the preschool pertussis booster vaccination: prospective cohort study. *Bmj*. 2014;348:g3668.
 25. Tubeuf S, Edlin R, Shourie S, Cheater FM, Bekker H, Jackson C. Cost effectiveness of a web-based decision aid for parents deciding about MMR vaccination: a three-arm cluster randomised controlled trial in primary care. *The British journal of general practice : the journal of the Royal College of General Practitioners*. 2014;64(625):e493-9.
 26. Said A, Jou JH. Hepatitis B vaccination and screening awareness in primary care practitioners. *Hepatitis research and treatment*. 2014;2014:373212.
 27. Nessler K, Krzton-Krolewiecka A, Chmielowiec T, Jarczewska D, Windak A. Determinants of influenza vaccination coverage rates among primary care patients in Krakow, Poland and the surrounding region. *Vaccine*. 2014;32(52):7122-7.
 28. Miranda Garcia M, Gutierrez Teira B, Ariza Cardiel G. [A study on the hepatitis B virus vaccination rates in hepatitis C-infected patients in a Primary Care centre]. *Atencion primaria*. 2014;46(7):395.
 29. Maurer J, Harris KM. Issuance of patient reminders for influenza vaccination by US-based primary care physicians during the first year of universal influenza vaccination recommendations. *American journal of public health*. 2014;104(6):e60-2.
 30. Liyanage H, de Lusignan S, Liaw ST, Kuziemyk CE, Mold F, Krause P, et al. Big Data Usage Patterns in the Health Care Domain: A Use Case Driven Approach Applied to the Assessment of Vaccination Benefits and Risks. *Contribution of the IMIA Primary Healthcare Working Group. Yearbook of medical informatics*. 2014;9:27-35.
 31. Haas-Gehres A, Sebastian S, Lamberjack K. Impact of pharmacist integration in a pediatric primary care clinic on vaccination errors: a retrospective review. *Journal of the American Pharmacists Association : JAPhA*. 2014;54(4):415-8.
 32. Fitchett JR, Arnott ND. Influenza vaccination uptake among people aged over 85 years: an audit of primary care practice in the UK. *JRSM open*. 2014;5(7):2054270414531122.

33. Fiscella K. Tackling disparities in influenza vaccination in primary care: it takes a team. *Journal of general internal medicine*. 2014;29(12):1579-81.
34. Bruno DM, Wilson TE, Gany F, Aragones A. Identifying human papillomavirus vaccination practices among primary care providers of minority, low-income and immigrant patient populations. *Vaccine*. 2014;32(33):4149-54.
35. Wong MC, Lee A, Ngai KL, Chor JC, Chan PK. Knowledge, attitude, practice and barriers on vaccination against human papillomavirus infection: a cross-sectional study among primary care physicians in Hong Kong. *PLoS one*. 2013;8(8):e71827.
36. Maltezou HC, Katerelos P, Poufta S, Pavli A, Maragos A, Theodoridou M. Attitudes toward mandatory occupational vaccinations and vaccination coverage against vaccine-preventable diseases of health care workers in primary health care centers. *American journal of infection control*. 2013;41(1):66-70.
37. da Costa FM, de Barros Lima Martins AM, Dos Santos Neto PE, de Pinho Veloso DN, Magalhaes VS, Ferreira RC. Is vaccination against hepatitis B a reality among primary health care workers? *Revista latino-americana de enfermagem*. 2013;21(1):316-24.
38. Chu D, Yang JD, Lok AS, Tran T, Martins EB, Fagan E, et al. Hepatitis B screening and vaccination practices in asian american primary care. *Gut and liver*. 2013;7(4):450-7.
39. Chaudhry R, Schieter SM, North F, Dejesus R, Kesman RL, Stroebel RJ. Improving rates of herpes zoster vaccination with a clinical decision support system in a primary care practice. *Journal of evaluation in clinical practice*. 2013;19(2):263-6.
40. Aziz NA, Muhamad S, Manaf MR, Hamid MZ. Factors Influencing H1N1 Vaccination Among Primary Health Care Workers: A Cross-Sectional Study. *International journal of preventive medicine*. 2013;4(6):664-70.
41. Allison MA, Dunne EF, Markowitz LE, O'Leary ST, Crane LA, Hurley LP, et al. HPV vaccination of boys in primary care practices. *Academic pediatrics*. 2013;13(5):466-74.
42. Simpson CR, McMenamin J. Pandemic influenza vaccination for healthcare workers in primary care: good progress, but higher uptake required. *Primary care respiratory journal : journal of the General Practice Airways Group*. 2012;21(3):246-7.
43. Real W, Alvarado V. [Vaccination against human papilloma virus: a Chilean experience at primary health care centers]. *Revista chilena de infectologia : organo oficial de la Sociedad Chilena de Infectologia*. 2012;29(6):686.
44. Pruvost I, Lowingerova-Fauvet S, Debacker P, Dervaux B, Dubos F, Martinot A. [Assessment of primary care physicians' adherence to the national childhood vaccination schedule]. *Archives de pediatrie : organe officiel de la Societe francaise de pediatrie*. 2012;19(3):248-53.
45. O'Leary ST, Stokley S, Crane LA, Allison MA, Hurley LP, Wortley P, et al. Influenza vaccination in the 2009-2010 pandemic season: the experience of primary care physicians. *Preventive medicine*. 2012;55(1):68-71.
46. Maceira Castineira MM, Deben Sanchez M, Ramil Hermida L, Acevedo Prado A. [Vaccination of primary health care staff in the area of Ferrol (Galicia) against seasonal and swine flu]. *Atencion primaria*. 2012;44(3):180-1.
47. Clark SJ, Lamarand K, Dombkowski KJ. Identifying children with chronic conditions for influenza vaccination using a statewide immunization registry: initial experiences of primary care providers. *Journal of public health management and practice : JPHMP*. 2012;18(3):204-8.
48. Banach DB, Ornstein K, Factor SH, Soriano TA. Seasonal influenza vaccination among homebound elderly receiving home-based primary care in New York City. *Journal of community health*. 2012;37(1):10-4.
49. Almazan Altuzarra J, Arana Alvarez K, Gonzalez Suarez JC, Ruiz Begue M. [Is influenza vaccination the new Bellerophon? The flu vaccine in primary care]. *Semergen*. 2012;38(3):197-8.
50. Albert SM, Nowalk MP, Yonas MA, Zimmerman RK, Ahmed F. Standing orders for influenza and pneumococcal polysaccharide vaccination: correlates identified in a national survey of U.S. Primary care physicians. *BMC family practice*. 2012;13:22.
51. Zimmerman RK, Albert SM, Nowalk MP, Yonas MA, Ahmed F. Use of standing orders for adult influenza vaccination: a national survey of primary care physicians. *American journal of preventive medicine*. 2011;40(2):144-8.